

# False Hope

The rate of global temperature rise may have hit a plateau, but a climate crisis still looms in the near future

By Michael E. Mann

“Temperatures have been flat for 15 years—nobody can properly explain it,” the *Wall Street Journal* says. “Global warming ‘pause’ may last for 20 more years, and Arctic sea ice has already started to recover,” the *Daily Mail* says. Such reassuring claims about climate abound in the popular media, but they are misleading at best. Global warming continues unabated, and it remains an urgent problem.

The misunderstanding stems from data showing that during the past decade there was a slowing in the rate at which the earth’s average surface temperature had been increasing. The event is commonly referred to as “the pause,” but that is a misnomer: temperatures still rose, just not as fast as during the prior decade. The important question is, What does the short-term slowdown portend for how the world may warm in the future?

The Intergovernmental Panel on Climate Change (IPCC) is charged with answering such questions. In response to the data, the IPCC in its September 2013 report lowered one aspect of its prediction for future warming. Its forecasts, released every five to seven years, drive climate policy worldwide, so even the small change raised debate over how fast the planet is warming and how much time we have to stop it. The IPCC has not yet weighed in on the impacts of the warming or how to mit-


igate it, which it will do in reports that were due this March and April. Yet I have done some calculations that I think can answer those questions now: If the world keeps burning fossil fuels at the current rate, it will cross a threshold into environmental ruin by 2036. The “faux pause” could buy the planet a few extra years beyond that date to reduce greenhouse gas emissions and avoid the crossover—but only a few.

### A SENSITIVE DEBATE

THE DRAMATIC NATURE of global warming captured world attention in 2001, when the IPCC published a graph that my co-authors and I devised, which became known as the “hockey stick.” The shaft of the stick, horizontal and sloping gently downward from left to right, indicated only modest changes in Northern Hemisphere temperature for almost 1,000 years—as far back as our data went.

### Danger Zone in 22 Years

If the Northern Hemisphere’s surface temperatures rise more than two degrees Celsius above preindustrial levels (*baseline*), human civilization will suffer dangerous harm, scientists say. When will that occur if the world keeps burning fossil fuels at current rates? The answer comes from entering estimates for equilibrium climate sensitivity (ECS)—how sensitive the atmosphere is to the heating effect of greenhouse gases (*five solid curves*)—into a so-called energy balance model of climate. The estimate that best agrees with recorded data reflecting the sensitivity of the earth’s climate (*white*) indicates that the world will cross the two degrees C threshold in 2036, only 22 years from now (*orange*). If the reported recent slowdown in the rate of temperature rise, sometimes inappropriately called “the pause,” proves to be part of a more persistent pattern, then a different estimate (*gold*) best fits the past 15 years or so, and it gives the world until 2046 to cross the danger line.



### Where to Hold the Line

Scientists and policy makers commonly say that the world has to keep atmospheric CO<sub>2</sub> levels below 450 ppm to avoid two degrees C of warming (the level briefly hit 400 ppm in 2013). Yet if the atmosphere’s climate sensitivity is three degrees C (*orange*), warming can be limited to that amount only if we keep emitting polluting aerosols (particles in the atmosphere that partly block the sun’s heat) at current rates (*dashed orange*). Ironically, the reduction in coal burning needed to lower CO<sub>2</sub> emissions also lessens aerosols, sending temperatures across the danger line (*dotted orange*). The same is true if the sensitivity is 2.5 degrees C (*gold*). These data therefore indicate that to reliably avoid two degrees C of warming, CO<sub>2</sub> levels should be held to 405 ppm (*blue*)—barely above the 393 to 400 ppm levels observed in the past year.

The upturned blade of the stick, at the right, indicated an abrupt and unprecedented rise since the mid-1800s. The graph became a lightning rod in the climate change debate, and I, as a result, reluctantly became a public figure. In its September 2013 report, the IPCC extended the stick back in time, concluding that the recent warming was likely unprecedented for at least 1,400 years.

Although the earth has experienced exceptional warming over the past century, to estimate how much more will occur we need to know how temperature will respond to the ongoing human-caused rise in atmospheric greenhouse gases, primarily carbon dioxide. Scientists call this responsiveness “equilibrium climate sensitivity” (ECS). ECS is a common measure of the heating effect of greenhouse gases. It represents the warming at the earth’s surface that is expected after the concentration of CO<sub>2</sub> in the atmosphere doubles and the climate subsequently stabilizes (reaches equilibrium).

The preindustrial level of CO<sub>2</sub> was about 280 parts per million (ppm), so double is roughly 560 ppm. Scientists expect this doubling to occur later this century if nations continue to burn fossil fuels as they do now—the “business as usual” scenario—instead of curtailing fossil-fuel use. The more sensitive the atmosphere is to a rise in CO<sub>2</sub>, the higher the ECS, and the faster the temperature will rise. ECS is shorthand for the amount of warming expected, given a particular fossil-fuel emissions scenario.

It is difficult to determine an exact value of ECS because warming is affected by feedback mechanisms, including clouds, ice and other factors. Different modeling groups come to different conclusions on what the precise effects of these feedbacks may be. Clouds could be the most significant. They can have both a cooling effect, by blocking out incoming sunlight, and a warming effect, by absorbing some of the heat energy that the earth sends out toward space. Which of these effects dominates depends on the type, distribution and altitude of the clouds—difficult for climate models to predict. Other feedback factors relate to how much water vapor there will be in a warmer atmosphere and how fast sea ice and continental ice sheets will melt.

Because the nature of these feedback factors is uncertain, the IPCC provides a range for ECS, rather than a single number. In the September report—the IPCC’s fifth major assessment—the panel settled on a range of 1.5 to 4.5 degrees Celsius (roughly three to eight degrees Fahrenheit). The IPCC had lowered the bottom end of the range, down from the two degrees C it had set in its Fourth Assessment Report, issued in 2007. The IPCC based the lowered bound on one narrow line of evidence: the slowing of surface warming during the past decade—yes, the faux pause.

Many climate scientists—myself included—think that a single decade is too brief to accurately measure global warming and that the IPCC was unduly influenced by this one, short-term number. Furthermore, other explanations for the speed bump do not contradict the preponderance of evidence that suggests that temperatures will continue to rise. For example, the accumulated effect of volcanic eruptions during the past decade, including

**Michael E. Mann** is Distinguished Professor of Meteorology at Pennsylvania State University and contributed to the International Panel on Climate Change work that received the 2007 Nobel Peace Prize. His book *The Hockey Stick and the Climate Wars: Dispatches from the Front Lines* (Columbia University Press, 2012) is in paperback, with a foreword by Bill Nye the Science Guy.



the Icelandic volcano with the impossible name, Eyjafjallajökull, may have had a greater cooling effect on the earth’s surface than has been accounted for in most climate model simulations. There was also a slight but measurable decrease in the sun’s output that was not taken into account in the IPCC’s simulations.

Natural variability in the amount of heat the oceans absorb may have played a role. In the latter half of the decade, La Niña conditions persisted in the eastern and central tropical Pacific, keeping global surface temperatures about 0.1 degree C colder than average—a small effect compared with long-term global warming but a substantial one over a decade. Finally, one recent study suggests that incomplete sampling of Arctic temperatures led to underestimation of how much the globe actually warmed.

None of these plausible explanations would imply that climate is less sensitive to greenhouse gases. Other measurements also do not support the IPCC’s revised lower bound of 1.5 degrees C. When all the forms of evidence are combined, they point to a most likely value for ECS that is close to three degrees C. And as it turns out, the climate models the IPCC actually used in its Fifth Assessment Report imply an even higher value of 3.2 degrees C. The IPCC’s lower bound for ECS, in other words, probably does not have much significance for future world climate—and neither does the faux pause.

For argument’s sake, however, let us take the pause at face value. What would it mean if the actual ECS were half a degree lower than previously thought? Would it change the risks presented by business-as-usual fossil-fuel burning? How quickly would the earth cross the critical threshold?

#### A DATE WITH DESTINY: 2036


MOST SCIENTISTS CONCUR that two degrees C of warming above the temperature during preindustrial time would harm all sectors of civilization—food, water, health, land, national security, energy and economic prosperity. ECS is a guide to when that will happen if we continue emitting CO<sub>2</sub> at our business-as-usual pace.

I recently calculated hypothetical future temperatures by plugging different ECS values into a so-called energy balance model, which scientists use to investigate possible climate scenarios. The computer model determines how the average surface temperature responds to changing natural factors, such as volcanoes and the sun, and human factors—greenhouse gases, aerosol pollutants, and so on. (Although climate models have critics, they reflect our best ability to describe how the climate system works, based on physics, chemistry and biology. And they have a proved track

#### PREDICTING THE FUTURE

### A Solid Line of Evidence

Determining when the planet’s atmosphere will cross the dangerous warming threshold of two degrees C [see graph on pages 78 and 79] depends on how sensitive the atmosphere is to rising CO<sub>2</sub> levels. The most likely value for this equilibrium climate sensitivity (*horizontal axis*) is just below three degrees C. Why? Because many independent calculations of temperature in the distant past, as well as many climate models, place the number very close to this value (*various color bars*). The product of all the lines of evidence appears at the bottom (*gray bar*).



SOURCE: “THE EQUILIBRIUM SENSITIVITY OF THE EARTH’S TEMPERATURE TO RADIATION CHANGES,” BY RETO KNUTTI AND GABRIELE C. HEGGERL, IN *NATURE GEOSCIENCE*, VOL. 1, NOVEMBER 2008

record: for example, the actual warming in recent years was accurately predicted by the models decades ago.)

I then instructed the model to project forward under the assumption of business-as-usual greenhouse gas emissions. I ran the model again and again, for ECS values ranging from the IPCC’s lower bound (1.5 degrees C) to its upper bound (4.5 degrees C). The curves for an ECS of 2.5 degrees and three degrees C fit the instrument readings most closely. The curves for a substantially lower (1.5 degrees C) and higher (4.5 degrees C) ECS did not fit the recent instrumental record at all, reinforcing the notion that they are not realistic.

To my wonder, I found that for an ECS of three degrees C, our planet would cross the dangerous warming threshold of two degrees C in 2036, only 22 years from now. When I considered the lower ECS value of 2.5 degrees C, the world would cross the threshold in 2046, just 10 years later [see graph on pages 78 and 79].

So even if we accept a lower ECS value, it hardly signals the end of global warming or even a pause. Instead it simply buys us a little bit of time—potentially valuable time—to prevent our planet from crossing the threshold.

#### CAUTIOUS OPTIMISM

THESE FINDINGS HAVE IMPLICATIONS for what we all must do to prevent disaster. An ECS of three degrees C means that if we are to limit global warming to below two degrees C forever, we need to keep CO<sub>2</sub> concentrations far below twice preindustrial levels, closer to 450 ppm. Ironically, if the world burns significantly less coal, that would lessen CO<sub>2</sub> emissions but also reduce aerosols in the atmosphere that block the sun (such as sulfate particulates), so we would have to limit CO<sub>2</sub> to below roughly 405 ppm.

We are well on our way to surpassing these limits. In 2013 atmospheric CO<sub>2</sub> briefly reached 400 ppm for the first time in recorded history—and perhaps for the first time in millions of years, according to geologic evidence. To avoid breaching the 405-ppm threshold, fossil-fuel burning would essentially have to cease immediately. To avoid the 450-ppm threshold, global carbon emissions could rise only for a few more years and then would have to ramp down by several percent a year. That is a tall task. If the ECS is indeed 2.5 degrees C, it will make that goal a bit easier.

Even so, there is considerable reason for concern. The conclusion that limiting CO<sub>2</sub> below 450 ppm will prevent warming beyond two degrees C is based on a conservative definition of climate sensitivity that considers only the so-called fast feedbacks in the climate system, such as changes in clouds, water vapor and melting sea ice. Some climate scientists, including James E. Hansen, former head of the NASA Goddard Institute for Space Studies, say we must also consider slower feedbacks such as changes in the continental ice sheets. When these are taken into account, Hansen and others maintain, we need to get back down to the lower level of CO<sub>2</sub> that existed during the mid-20th century—about 350 ppm. That would require widespread deployment of expensive “air capture” technology that actively removes CO<sub>2</sub> from the atmosphere.

Furthermore, the notion that two degrees C of warming is a “safe” limit is subjective. It is based on when *most* of the globe will be exposed to potentially irreversible climate changes. Yet destructive change has already arrived in some regions. In the Arctic, loss of sea ice and thawing permafrost are wreaking havoc on indigenous peoples and ecosystems. In low-lying island nations, land and freshwater are disappearing because of rising sea levels and erosion. For these regions, current warming, and the further warming (at least 0.5 degree C) guaranteed by CO<sub>2</sub> already emitted, constitutes damaging climate change today.

Let us hope that a lower climate sensitivity of 2.5 degrees C turns out to be correct. If so, it offers cautious optimism. It provides encouragement that we can avert irreparable harm to our planet. That is, if—and only if—we accept the urgency of making a transition away from our reliance on fossil fuels for energy. ■

#### MORE TO EXPLORE

**Defining Dangerous Anthropogenic Interference.** Michael E. Mann in *Proceedings of the National Academy of Sciences USA*, Vol. 106, No. 11, pages 4065–4066; March 17, 2009.

**Separating Forced from Chaotic Climate Variability over the Past Millennium.** Andrew P. Schurer et al. in *Journal of Climate*, Vol. 26, No. 18, pages 6954–6973; September 2013.

#### FROM OUR ARCHIVES

**Global Warming: Faster Than Expected?** John Carey; November 2012.

#### IN BRIEF

**The rate** at which the earth’s temperature has been rising eased slightly in the past decade, but temperature is still increasing; calling the slowdown a “pause” is false.

**New calculations** by the author indicate that if the world continues to burn fossil fuels at the current rate, global warming will rise to two degrees Celsius by 2036,

crossing a threshold that will harm human civilization. **To avoid** the threshold, nations will have to keep carbon dioxide levels below 405 parts per million.